

Molecular Survey of Zoonotic Agents in Rodents and Other Small Mammals in Croatia

Ante Tadin,† Rafal Tokarz,*† Alemka Markotić, Josip Margaletić, Nenad Turk, Josipa Habuš, Petra Svoboda, Marko Vucelja, Aaloki Desai, Komal Jain, and W. Ian Lipkin

University Hospital for Infectious Diseases “Dr. Fran Mihaljević,” Zagreb, Croatia; Center for Infection and Immunity, Mailman School of Public Health, Columbia University, New York, New York; Faculty of Forestry, University of Zagreb, Zagreb, Croatia; Faculty of Veterinary Medicine, University of Zagreb, Zagreb, Croatia

Abstract. Croatia is a focus for many rodent-borne zoonosis. Here, we report a survey of 242 rodents and small mammals, including 43 *Myodes glareolus*, 131 *Apodemus flavicollis*, 53 *Apodemus agrarius*, three *Apodemus sylvaticus*, six *Sorex araneus*, four *Microtus arvalis*, one *Microtus agrestis*, and one *Muscardinus avellanarius*, collected at eight sites in Croatia over an 8-year period. Multiplex MassTag polymerase chain reaction (PCR) was used for detection of *Borrelia*, *Rickettsia*, *Bartonella*, *Babesia*, *Ehrlichia*, *Anaplasma*, *Francisella tularensis*, and *Coxiella burnetii*. Individual PCR assays were used for detection of *Leptospira*, lymphocytic choriomeningitis virus, orthopoxviruses, flaviviruses, hantaviruses, and *Toxoplasma gondii*. Of the rodents, 52 (21.5%) were infected with *Leptospira*, 9 (3.7%) with *Borrelia miyamotoi*, 5 (2%) with *Borrelia afzelii*, 29 (12.0%) with *Bartonella*, 8 (3.3%) with *Babesia microti*, 2 (0.8%) with *Ehrlichia*, 4 (1.7%) with *Anaplasma*, 2 (0.8%) with *F. tularensis*, 43 (17.8%) with hantaviruses, and 1 (0.4%) with an orthopoxvirus. Other agents were not detected. Multiple infections were found in 32 rodents (13.2%): dual infections in 26 rodents (10.7%), triple infections in four rodents (2.9%), and quadruple infections in two rodents (0.8%). Our findings indicate that rodents in Croatia harbor a wide range of bacteria and viruses that are pathogenic to humans.

INTRODUCTION

Rodents are hosts to a wide range of zoonotic pathogens. Rodent-borne agents can be transmitted to humans directly through contact with rodents and their excretions or through rodent bites and indirectly by arthropod vectors including ticks, fleas, and mites. Croatia is a natural focus for many rodent-borne zoonoses because of its diverse forest ecology and wide range of rodent species. Different rodent species in Croatia have been implicated as reservoirs to several clinically important bacteria, protozoa, and viruses. The main rodent-associated bacteria include species of *Anaplasma*, *Borrelia*, *Bartonella*, *Coxiella*, *Ehrlichia*, *Francisella*, *Leptospira*, and *Rickettsia*. Infections with *Leptospira* and *Borrelia* are the most frequently reported from among this group. With a mean yearly infection incidence of 1.7 per 100,000 inhabitants, Croatia has one of the highest recorded incidences of human leptospirosis in the world.¹ *Borrelia* infections result in two distinct tick-transmitted diseases, Lyme borreliosis and relapsing fever (RF). Both are endemic in Croatia with majority of cases of borreliosis reported in the northwest of the country.² Q fever, caused by *Coxiella burnetii* is endemic in parts of southern Croatia.^{3,4} Limited data exist on tick-transmitted *Rickettsia*, *Anaplasma*, and *Ehrlichia*.⁵ Human cases of anaplasmosis and rickettsiosis have been reported in Croatia, but little data are available on ehrlichiosis.^{6,7} Infections with *Francisella tularensis*, the agent of tularemia, are occasionally reported in Croatia, particularly in Sava Valley in the central part of the country.⁸ *Bartonella* spp. are hemotropic bacteria detected in many mammals, predominantly rodents.⁹ Many species and subspecies of *Bartonella* circulate in European rodents, although the pathogenicity of many of these species is as yet undetermined. In addition, the diver-

sity of *Bartonella* spp. present in rodents in Croatia has not been examined.

Hantaviruses are the viruses most frequently associated with rodents in Croatia. These include Dobrava (DOBV) and Puumala (PUUV) viruses, implicated in hemorrhagic fever with renal syndrome, and the nonpathogenic Tula (TULV) and Saaremaa viruses.^{10–14} Rodents can also be infected with flaviviruses, orthopoxviruses, and lymphocytic choriomeningitis virus (LCMV); however, to our knowledge, there are no published reports of investigation for the presence of these viruses in Croatia.^{15–17} Rodents in Croatia are also reservoirs of protozoan *Apicomplexa* parasites, including *Toxoplasma* and *Babesia* that cause toxoplasmosis and babesiosis, respectively, mainly in immunocompromised individuals.

Here, we report results of analysis of rodents and small mammals trapped in eight localities in Croatia for the presence of a wide range of known zoonotic human pathogens.

MATERIALS AND METHODS

Sample collection. This survey was an extension of a project designed to assess the distribution of small rodents in diverse forest ecosystems in Croatia. Between 2003 and 2011, rodents were trapped in eight different localities in Croatia: Ivanić-Grad, Mikanovci, Ilok, Cerna, Papuk, Žutica, Draganić, and Sušica (Figure 1). In each of three selected geographic transects at each location, 100 traps were placed. The distance between the traps was approximately 7 m and the distance between the transects was approximately 150 m. Each transect was at least 50 m away from the edge of the forest stands. Each of these localities includes forest ecosystems without proximity to urban areas. All rodents were trapped using Sherman-type live traps and were morphologically characterized to species.¹⁸ Species identification of *Apodemus flavicollis* and *Apodemus sylvaticus* was confirmed by PCR (polymerase chain reaction) targeting the mitochondrial cytochrome *b* gene and sequencing of PCR products.^{19,20} A total of 242 animals were collected, including 131 yellow-necked mouse (*A. flavicollis*), 53 striped field mouse (*Apodemus agrarius*), three wood mouse (*A. sylvaticus*),

* Address correspondence to Rafal Tokarz, Center for Infection and Immunity, Mailman School of Public Health, Columbia University, 722 West 168th Street, New York, NY 10032. E-mail: rt2249@cumc.columbia.edu

† These authors contributed equally to this work.



FIGURE 1. Map in Croatia indicating the regions where the rodents were trapped.

43 bank voles (*Myodes glareolus*), six common shrews (*Sorex araneus*), four common voles (*Microtus arvalis*), one field vole (*Microtus agrestis*), and one common dormouse (*Muscardinus avellanarius*). Animal experimentation guidelines of the American Society of Mammologists was followed during all animal work.²¹ After euthanasia, kidney and lung samples were collected from individual specimens and stored at -80°C .

Nucleic acid extraction and complementary DNA synthesis. A total of 323 organ specimens were available for PCR analysis. These consisted of lung samples acquired from 237 animals and kidney samples from 86 animals. At least one organ (lung or kidney) was available for PCR analysis from all 242 animals. In cases where both lung and kidney samples were available from an individual animal, both were tested by PCR. Tissues were homogenized using TissueLyser and 5-mm beads for homogenization (Qiagen, Hilden, Germany). DNA and RNA were extracted from each sample using the AllPrep DNA/RNA Mini Kit (Qiagen). DNA from each sample was eluted in 100 μL EB Buffer and RNA from each sample was eluted in 40 μL RNase-free water. Nucleic acid concentration was assessed with the NanoDrop Spectrophotometer (Thermo Scientific, Waltham, MA). Complementary DNA (cDNA) was prepared in a 20 μL reaction using 5 μL extracted RNA and SuperScript II Reverse Transcriptase (Invitrogen, Carlsbad, CA).

Polymerase chain reaction. MassTag PCR was performed using primers for detection of Lyme borreliosis-associated *Borrelia* spp., *Bartonella* spp., *Ehrlichia* spp., *Anaplasma* spp.,

F. tularensis, *C. burnetii*, *Rickettsia* spp., RF *Borrelia* spp., and *Babesia* spp. (Table 1).^{22,23,29} Of template DNA, 2 μL was used in all reactions. Reaction conditions were 94°C for 15 minutes; one cycle at 94°C for 30 seconds, 65°C for 30 seconds, and 72°C for 30 seconds, followed by 11 cycles with annealing temperature decreased by 1°C in each cycle. The final PCR was run for 38 cycles at 94°C for 30 seconds, 54°C for 30 seconds, and 72°C for 30 seconds.

All samples that were positive on the MassTag PCR assay were retested in individual PCR assays for the microbe(s) of interest using the same primer pairs. The presence of relevant sequence in the resulting PCR product was required as confirmation before a sample was declared agent positive. In cases where species differentiation was required, we designed primers within a discriminatory region of the genome to amplify partial sequences that were then compared with existing genomic data from GenBank. To discriminate among *Bartonella* spp., we designed primer pairs that amplified 513 nucleotides (nts) of the *gltA* gene and 800 nts of the *16SrRNA* gene. For *Babesia* spp., we used a primer pair that amplified 607 nts of the *18SrRNA* gene, and for *Borrelia* spp., an assay targeting the *flaB* gene amplified a sequence of 966 nts (Table 1). Sequences for *Bartonella* spp. and *Borrelia miyamotoi* were deposited in GenBank under accession nos. KT452901–KT452935. All alignments were generated in Geneious v 6.1.5 (Biomatters Ltd., Auckland, New Zealand). Maximum-likelihood phylogenetic tree for *Bartonella* was constructed using the Jukes–Cantor model with 1,000 bootstrap replicates in Mega 6.0.³⁰

TABLE 1
Primer sequences used for pathogen detection

Pathogen	Gene target	Primer sequence	Reference
Lyme borreliosis-associated <i>Borrelia</i> spp.	<i>flaB</i>	Fwd: GCAATGACAAAACATATTGRGGAATTGA Rev: YACAATGACMGATGAGGTTGTRGC	22
Relapsing fever <i>Borrelia</i> spp.	<i>flaB</i>	Fwd: GCTGAAGAGCTTGAATGCAAC Rev: GCAATTGCYTCATCCTGATTTG	23
<i>Bartonella</i> spp.	<i>hbpA</i>	Fwd: CTTCTGCRGCACAAGCTGCTGAT Rev: CCACCAATATARAAACCTGTCCAAGA	22
<i>Ehrlichia</i> spp.	<i>16SrRNA</i>	Fwd: CGTAAAGGGCACGTAAGTGGACTA Rev: CACCTCAGTGTCAGTATCGARCCA	22
<i>Anaplasma</i> spp.	<i>16SrRNA</i>	Fwd: GGGCATGTAGGCGGTTCCGGT Rev: TCAGCGTCAGTACCGGACCA	22
<i>Francisella tularensis</i>	<i>fopA</i>	Fwd: ATGTTTCGGCATGTGAATAGTTAA Rev: ACCACTGCTTTGTGTAGTAGCTGAA	22
<i>Coxiella burnetii</i>	<i>IS1111</i>	Fwd: GCTCCTCCACGCTTCCAT Rev: GGTTCAACTGTGTGGAATTGATGAGT	22
<i>Rickettsia</i> spp.	<i>ompB</i>	Fwd: YAAAAGTCAGCGTTACTTCTTYGA Rev: YGCTTRTTTGAACCTGTTGTACC	22
<i>Babesia</i> spp.	<i>18SrRNA</i>	Fwd: CGACTAGDGATTGGDGGTCGTC Rev: CHTGTCTGGACCTGGTGAGKTT	This study
Lymphocytic choriomeningitis virus	<i>RdRp</i>	Fwd: CCACTYTTGTCTGCACTGTCTAT Rev: CTTTTTGATGCGCAATGGAT	This study
Orthopoxvirus	<i>VETFS</i>	Fwd: ACCAACTATATTACCTCATCAGTTA Rev: TTAAACAAGTTCATAGCTACACCCA	This study
Flavivirus	<i>NS5</i>	Fwd: ATGGCHATGACNGACACNAC Rev: TTCTTYTCTCTYTTNCCCATCAT	This study
<i>Toxoplasma gondii</i>	<i>B1</i>	Fwd: GAAGAGATCCAGCAGATCTCGT Rev: TGAGAGGAGGCAGCACAAG	This study
Puumala/Tula hantavirus	<i>N</i>	Fwd1: TATGGIAATGTCCTTGATGT Rev1: GCACAIGCAAAAACCCA	24
		Fwd2: CCIAGTGGICAACAGC Rev2: AAICCIATIAICCCAT	24
Dobrava hantavirus	<i>G</i>	Fwd1: GGACCAGGTGCAGCTTGTGAAGC Rev1: ACCTCACAAACCATTGAACC	25
		Fwd2: ATGCCAGCGAGTCGACCAA Rev2: GAGCTATTATGTAAGATTGC	26
<i>Leptospira</i> spp.	<i>LA0322</i>	FwdA: CATTTCATGTTTCGAATCATTTCAA RevA: GCCCAAGTTCCTTCTAAAAG	27
	<i>secY</i>	FwdB: GAATTTCTCTTTTGATCTTCG RevB: GAGTTAGAGCTCAAATCTAAG	28
<i>Bartonella</i> spp.*	<i>gltA</i>	Fwd: GGTCTATCAYGACTCKATTGATATTA Rev: GCACGTGGRTCATAATTTTTATARAC	This study
	<i>16SrRNA</i>	Fwd: ATAGCAAGTCGAGCGCRCT Rev: GCCCGACGGCTAACATTC	This study
<i>Borrelia</i> spp.*	<i>flaB</i>	Fwd: GGGGATGATYATMAATCATAATAC Rev: TGCAATCATWGCCATTGC	This study
<i>Babesia</i> spp.*	<i>18SrRNA</i>	Fwd: TTCGACGGTAKGGTATTGGC Rev: GGGAATTTACCTCTGACAGTYAA	This study

*Consensus genus primers used for species discrimination.

All samples were additionally tested for LCMV, orthopoxvirus, flaviviruses, hantaviruses, and *Toxoplasma gondii* in single-agent PCR assays. For detection of RNA viruses, 2 µL cDNA were used as template. For the detection of orthopoxviruses and *T. gondii*, 2 µL DNA was used. Nested PCR assays were used to detect the presence of hantaviruses (PUUV, DOBV, and TULV).²⁴⁻²⁶ All primer sequences are listed in Table 1. PCR products were size-fractionated in 1% agarose gels and visualized using ethidium bromide before purification for dideoxy sequencing.

Detection of *Leptospira* by renal tissue culture and PCR. Kidney tissue from each animal ($N = 242$) was available for *Leptospira* detection. Tissue was inoculated into Korthof's medium for isolation and stabilization of *Leptospira* isolates. Isolates were grown to a density of $2-4 \times 10^8$ /mL and harvested by centrifugation at 14,000 rpm. Genomic DNA was extracted from each kidney tissue using the QIAamp DNA Mini Kit (Qiagen). Two PCR assays were used for species identification.

In the first assay, real-time PCR was used as described by Merien and others²⁷ with a primer set that amplified 331 nts of the locus LA0322 obtained from the complete genome sequence of *Leptospira interrogans* serovar Lai. All specimens were tested 10 times and mean melting temperatures were calculated to distinguish *Leptospira* genomic species. In a second assay, PCR was done with primers that amplified a 658-nt segment of *secY* that includes a 245-nt fragment suitable for rapid phylogenetic determination.²⁸ Amplified products were visualized in a 1.5% agarose gel, and their sizes were estimated by comparison with a 100-bp ladder (Amersham Biosciences, Little Chalfont, Buckinghamshire, United Kingdom). PCR products were purified by QIAquick PCR Purification Kit (Qiagen) and sequenced. DNA sequence clustal alignments were done using the LaserGene software package (DNASTAR, LaserGene, Madison, WI). In addition, a primer set that amplified a 563-nt DNA fragment of *Leptospira kirschneri* was used to confirm the speciation of *L. kirschneri* isolates.³¹

RESULTS

In our survey, 114 of 242 rodents (47%) were positive for at least one agent. Polymicrobial infections were detected in 32 rodents (13%). Dual infections were detected in 26 rodents (11%), triple infections in four rodents (3%), and quadruple infections in two rodents (0.8%) (Table 2).

Leptospira. *Leptospira* species were the most frequently detected agents with 52 *Leptospira*-positive rodents (21%). *Apodemus* spp. have previously been implicated as reservoirs of *Leptospira* in Croatia.^{32,33} Consistent with this earlier work, we found that 49 of 52 *Leptospira*-positive rodents represented *Apodemus* spp. Bacterial isolation from kidney tissue was successful in 40 PCR-positive rodents, and speciation assays indicated that 22 were *L. interrogans*, 16 *L. kirschneri*, and two *Leptospira borgpetersenii*.

Borrelia. Five rodents were positive for Lyme borreliosis-associated *Borrelia* spp. All were identified as *Borrelia afzelii* (> 99% identity with *B. afzelii* PKo strain, accession no. CP002933). The presence of this species in Croatian rodents has been reported previously.³⁴ Although *Borrelia garinii* is known to circulate in Croatia, we did not detect any rodents infected with this species. RF *Borrelia* was detected in nine rodents (4%) from four different localities. All RF *Borrelia* were identified as *B. miyamotoi* (> 99% identity with LB-2001 strain, accession no. CP006647) with the majority (7/9)

detected in *A. agrarius*. Two *B. miyamotoi*-positive rodents originated from Ivanić-Grad in the central part of Croatia, and the remaining seven were trapped in the eastern end of the country.

Bartonella. *Bartonella* spp. were detected in 29 animals (12%) in six different localities with the highest prevalence in Ilok and Draganić (26.7% and 25.0%, respectively). Although *A. flavicollis* was the most frequent reservoir of *Bartonella* in this study (9 of 29 *Bartonella*-positive rodents) and in earlier studies,³⁵ *A. agrarius* as well as *M. glareolus* and *S. araneus* were also positive for *Bartonella*. To determine the phylogeny of *Bartonella* in these rodents, we obtained partial sequences of *16SrRNA* and *gltA* genes by PCR. Previous reports have indicated that sequence analysis of *16SrRNA* gene alone is inadequate for species/genotype demarcation in *Bartonella*, and an additional loci need to be analyzed, with *gltA* the preferred gene used to establish *Bartonella* phylogeny.³⁶ Our analysis based on a 388-nt *gltA* sequence indicated that representatives of six putative *Bartonella* species were present in these rodents (Figure 2). Five rodents were infected with strains representative of *Bartonella grahamii* and nine rodents were infected with strains of *Bartonella taylorii*. Both species are distributed throughout Eurasia and are frequently detected in *Apodemus* and *Myodes* rodents.^{37,38} A single *A. flavicollis* was infected with a strain that clustered with *Bartonella elizabethae* species

TABLE 2
Summary of all agents and coinfections detected in different rodent species

	N	<i>Apodemus flavicollis</i>	<i>Apodemus agrarius</i>	<i>Apodemus sylvaticus</i>	<i>Myodes glareolus</i>	<i>Sorex araneus</i>	<i>Microtus arvalis</i>	<i>Microtus agrestis</i>	<i>Muscardinus avellanarius</i>
Total	242	131	53	3	43	6	4	1	1
<i>Borrelia afzelii</i>	5	2	1	–	3	–	–	–	–
<i>Bartonella</i> spp.	29	16	7	–	5	1	–	–	–
<i>Babesia microti</i>	8	4	3	–	–	–	–	1	–
<i>Ehrlichia</i> spp.	2	1	1	–	–	–	–	–	–
<i>Borrelia miyamotoi</i>	9	1	7	–	–	1	–	–	–
<i>Anaplasma</i> spp.	4	3	–	–	1	–	–	–	–
<i>Francisella tularensis</i>	2	–	2	–	–	–	–	–	–
<i>Leptospira</i> spp.	52	37	12	–	1	–	1	–	1
Hantavirus	43	36	2	–	4	–	1	–	–
DOBV	38	36	2	–	–	–	–	–	–
PUUV	4	–	–	–	4	–	–	–	–
TULV	1	–	–	–	–	–	1	–	–
Orthopoxvirus	1	–	–	–	1	–	–	–	–
Infections with two agents									
<i>Leptospira</i> , <i>B. afzelii</i>	1	1	–	–	–	–	–	–	–
<i>Leptospira</i> , <i>Ehrlichia</i>	1	–	1	–	–	–	–	–	–
<i>Leptospira</i> , <i>B. microti</i>	1	1	–	–	–	–	–	–	–
<i>Leptospira</i> , <i>Bartonella</i>	3	3	–	–	–	–	–	–	–
<i>Leptospira</i> , <i>Anaplasma</i>	1	1	–	–	–	–	–	–	–
<i>Leptospira</i> , <i>B. miyamotoi</i>	3	1	2	–	–	–	–	–	–
<i>Leptospira</i> , TULV	1	–	–	–	–	–	1	–	–
<i>Leptospira</i> , DOBV	10	10	–	–	–	–	–	–	–
<i>Bartonella</i> , DOBV	3	3	–	–	–	–	–	–	–
<i>B. microti</i> , <i>B. miyamotoi</i>	1	–	1	–	–	–	–	–	–
<i>Anaplasma</i> , DOBV	1	1	–	–	–	–	–	–	–
Infections with three agents									
<i>Leptospira</i> , <i>B. afzelii</i> , <i>B. microti</i>	1	–	1	–	–	–	–	–	–
<i>Bartonella</i> , <i>Anaplasma</i> , DOBV	1	1	–	–	–	–	–	–	–
<i>B. microti</i> , <i>Leptospira</i> , DOBV	2	2	–	–	–	–	–	–	–
Infections with four agents									
<i>Bartonella</i> , <i>B. afzelii</i> , <i>Leptospira</i> , DOBV	1	1	–	–	–	–	–	–	–
<i>Bartonella</i> , <i>F. tularensis</i> , <i>Leptospira</i> , DOBV	1	–	1	–	–	–	–	–	–

DOBV = Dobrava virus; PUUV = Puumala virus; TULV = Tula virus.




FIGURE 2. Maximum likelihood phylogenetic tree based on a 388-nucleotide fragment of *Bartonella gltA* gene. All sequences obtained in this study are indicated by * and the six species they cluster into are shown in gray. Only select relevant sequences from GenBank are shown in the tree.

group. Seven *A. flavicollis* were infected with a genotype similar to strain N40 isolated from *A. sylvaticus* from the United Kingdom.³⁹ On the basis of high sequence divergence relative to other strains and a distinct *gltA* phylogenetic clade formed by this group, we speculate that this genotype constitutes a new *Bartonella* species. Sequences obtained from six *A. agrarius* formed a monophyletic clade and likely constitute another potential novel *Bartonella* species. We also detected a single *Bartonella* strain (ARH567) in *A. agrarius* that clustered with AR15-3 strain from the United States (FN645480) and strains from Sweden (AY454539), Siberia (EF682090). Analysis of the *16SrRNA* sequence indicated that ARH567 was > 99% identical to *B. rochalimae* (strain BAA-1498, accession no. FN645459) and thus may constitute a *B. rochalimae* subspecies.

Babesia. Eight rodents from five localities were infected with *Babesia*, all identified as *Babesia microti* (100% identity to strain Jena, accession no. EF413181). Previous work has implicated *A. flavicollis* and *M. glareolus* as the primary reservoirs for *B. microti* in Europe.⁴⁰ Of the eight positive rodents in our study, three were *A. agrarius* and one *M. agrestis*, implicating these species as additional reservoirs of *Babesia* spp.

Hantaviruses. Hantaviruses were the second most frequently detected pathogens. Of 43 hantavirus-positive rodents, 38 were infected with DOBV, four with PUUV, and a single rodent was infected with TULV. Hantavirus-positive rodents originated from seven different locations with the highest prevalence in Žutica (34.5%) and Ilok (20%) (Table 3). Among *Apodemus* species, 28% of *A. flavicollis* and 2% of *A. agrarius* were DOBV positive. The highest prevalence of DOBV-positive rodents (30%) was in Žutica (Table 3). PUUV was detected in *M. glareolus* and TULV was detected in *M. arvalis*, similar to previous reports.¹²

Other agents. *Francisella tularensis* was detected in two *A. agrarius* from the Draganić site. One rodent was positive for orthopoxvirus. This is the first report of *F. tularensis* and orthopoxvirus infection in rodents in Croatia. Four rodents were positive for *Anaplasma* and two for *Ehrlichia*. *Rickettsia*, *C. burnetii*, LCMV, flaviviruses, and *T. gondii* were not detected. Among the agents detected, RF *Borrelia*, *Bartonella*, *Babesia*, *Ehrlichia*, and hantaviruses were detected in nucleic acids isolated from both lung and kidney tissues; *Anaplasma*, *Francisella*, and *B. afzelii* were detected in nucleic acids isolated only from lung tissue.

DISCUSSION

In this study, we demonstrate the presence of multiple zoonotic pathogens in rodent species present in Croatia. Although this study was not designed to demonstrate a linkage between the presence of these agents and human disease, the data presented here may provide insight into assessing the risk of rodent-borne zoonoses in Croatia.

Leptospira spp. and hantaviruses were the human pathogens most frequently detected in our study. The high prevalence of *Leptospira*-positive rodents parallels recent surveys in Croatia that revealed *Leptospira* infection in > 20% of rodents. The majority of *Leptospira*-positive rodents reported here were *Apodemus* species. In our previous work, *Leptospira* spp. were isolated from 13 of 28 *A. flavicollis* captured in Žutica forest.⁴¹ *Leptospira interrogans*, *L. kirschneri*, and *L. borgpetersenii* were identified as the *Leptospira* species in our study, similar to

TABLE 3
Number of agent-positive rodents at each trapping site

	Ivanić-Grad N = 39	Mikanovci N = 28	Ilok N = 30	Cerna N = 9	Papuk N = 18	Žutica N = 87	Draganić N = 16	Sušica N = 15	Total 242
<i>Borrelia afzelii</i>	–	1	–	–	–	3	0	1	5
<i>Bartonella</i>	4	1	8	–	2	10	4	–	29
<i>Babesia microti</i>	1	2	–	1	–	3	1	–	8
<i>Ehrlichia</i>	1	1	–	–	–	–	–	–	2
<i>Borrelia miyamotoi</i>	2	4	1	2	–	–	–	–	9
<i>Anaplasma</i>	–	–	–	–	3	1	–	–	4
<i>Francisella tularensis</i>	–	–	–	–	0	0	2	–	2
<i>Leptospira</i>	4	11	9	1	1	24	–	1	52
Hantaviruses	1	1	6	–	3	30	1	1	43
DOBV	1	–	6	–	3	26	1	1	38
PUUV	–	–	–	–	–	4	–	–	4
TULV	–	1	–	–	–	–	–	–	1
Orthopoxvirus	–	–	–	–	–	1	–	–	1

DOBV = Dobrava virus; PUUV = Puumala virus; TULV = Tula virus.

previous reports from Croatian rodents.^{32,33} Hantavirus and *Leptospira* also represented the most frequent coinfections. We previously showed coinfections with both agents in humans¹⁰ and rodents^{41,42} highlighting the potential risk for simultaneous transmission of these pathogens. The clinical underestimation of hantavirus and *Leptospira* coinfections was recently emphasized.⁴³

The genus *Bartonella* contains over 30 named species. Approximately 20 are adapted to wild rodents and six of these have caused documented human infections.³⁷ We detected the presence of six putative species present in Croatian rodents. Our data provide insight into the diversity of *Bartonella* in Croatia and the high overall genetic diversity of this genus worldwide. The extent of human infections associated with rodent-borne *Bartonella* is unknown as data regarding the incidence of zoonotic human *Bartonella* infections in Europe are not available. However, despite a relatively high prevalence of infected rodents, based on existing reports of infections due to rodent-borne *Bartonella*, we anticipate that the risk of human infection is low.

In this study, we report the first evidence of *Borrelia miyamotoi* in Croatia and have implicated *Apodemus* mice as reservoirs, similar to recent report from Switzerland.⁴⁴ *Borrelia miyamotoi* is an emerging tick-borne pathogen phylogenetically classified within the RF *Borrelia* group. Originally detected in Asia, it was subsequently detected in North America and Europe.^{23,45,46} Recent work has also implicated it as a potential cause of meningoencephalitis.^{47,48}

The prevalence of *Anaplasma* spp. infection in our study was 1.7%. *Anaplasma* spp. were detected in only two locations in central Croatia (Papuk and Žutica) among *A. flavicollis* and *M. glareolus*. *Anaplasma* spp. were previously reported in small rodents elsewhere in Europe,⁴⁹ but not in Croatia. *Ehrlichia* spp. were detected in only two rodents (0.8%) of genus *Apodemus* in two locations (Ivanić-Grad and Mikanovci).

A limitation of this study was the unavailability of organs other than lungs and kidneys for analysis. This is because we used samples banked during the course of a hantavirus and *Leptospira* surveillance project wherein only lungs and kidneys were collected. Thus, we may have missed some bacteria and viruses that concentrate in other organs. Furthermore, we did not screen for all rodent-associated agents, nor did we

pursue unbiased sequencing. Nonetheless, our data indicate that small rodents in Croatia are reservoirs for a wide range of viral, bacterial, and protozoan pathogens. We also demonstrated a high rate of coinfections that may have important implications for simultaneous transmission to humans.

Received July 10, 2015. Accepted for publication November 2, 2015.

Published online December 28, 2015.

Financial support: Work reported here was supported by awards from the NIH (AI057158, Northeast Biodefense Center-Lipkin), USAID-PREDICT, DTRA, and by the grants 143-1430115-0103, 068-1430115-2119, 053-1430115-2116 from the Ministry of Science, Education and Sports of Republic Croatia and the project “The role of biotic agents on vitality of narrow-leaved ash in Croatian floodplain forests” from Croatian Science Foundation.

Authors' addresses: Ante Tadin, Alemka Markotić, and Petra Svoboda, University Hospital for Infectious Diseases “Dr. Fran Mihaljević,” Zagreb, Croatia, E-mails: ante.tadin@yahoo.com, alemka.markotic@gmail.com, and petra.petrsvoboda@gmail.com. Rafal Tokarz, Aaloki Desai, Komal Jain, and W. Ian Lipkin, Center for Infection and Immunity, Mailman School of Public Health, Columbia University, New York, NY, E-mails: rt2249@cumc.columbia.edu, as3745@cumc.columbia.edu, kj2230@cumc.columbia.edu, and wil2001@cumc.columbia.edu. Josip Margaletić and Marko Vucelja, Faculty of Forestry, University of Zagreb, Zagreb, Croatia, E-mails: josip.margaletic@sumfak.hr and vucelja.marko@gmail.com. Nenad Turk and Josipa Habuš, Faculty of Veterinary Medicine, University of Zagreb, Zagreb, Croatia, E-mails: turk@vef.hr. and jhabus@vef.hr.

REFERENCES

1. Topic MB, Habus J, Milas Z, Tosev EC, Stritof Z, Turk N, 2010. Human leptospirosis in Croatia: current status of epidemiology and clinical characteristics. *Trans R Soc Trop Med Hyg* 104: 202–206.
2. Mulic R, Antonijevic S, Klismanic Z, Ropac D, Lucev O, 2006. Epidemiological characteristics and clinical manifestations of Lyme borreliosis in Croatia. *Mil Med* 171: 1105–1109.
3. Milotic I, Miletic B, Morovic M, 2001. Clinical, epidemiological and epizootic features of Q fever in the northern coastal part of Croatia from 1989 to 1998. *Acta Med Croatica* 55: 53–57.
4. Arricau-Bouvery N, Rodolakis A, 2005. Is Q fever an emerging or re-emerging zoonosis? *Vet Res* 36: 327–349.
5. Schex S, Dobler G, Riehm J, Muller J, Essbauer S, 2011. *Rickettsia* spp. in wild small mammals in Lower Bavaria, south-eastern Germany. *Vector Borne Zoonotic Dis* 11: 493–502.
6. Punda-Polic V, Luksic B, Capkun V, 2008. Epidemiological features of Mediterranean spotted fever, murine typhus, and Q

- fever in Split-Dalmatia County (Croatia), 1982–2002. *Epidemiol Infect* 136: 972–979.
7. Misić-Majeruš L, Bujic N, Madarić V, Avsić-Zupanc T, Milinković S, 2006. Human anaplasmosis (ehrlichiosis) [in Croatian]. *Acta Med Croatica* 60: 411–419.
 8. Borčić B, Hrabar A, Dulic B, Tvrtković N, Bilic V, Mikacic D, 1976. Ecological features of the tularemia natural focus in central Posavina (Croatia). *Folia Parasitol (Praha)* 23: 257–265.
 9. Kosoy M, Hayman DT, Chan KS, 2012. Bartonella bacteria in nature: where does population variability end and a species start? *Infect Genet Evol* 12: 894–904.
 10. Markotić A, Kuzman I, Babić K, Gagro A, Nichol S, Ksiazek TG, Rabatić S, Dekarić D, 2002. Double trouble: hemorrhagic fever with renal syndrome and leptospirosis. *Scand J Infect Dis* 34: 221–224.
 11. Markotić A, Nichol ST, Kuzman I, Sanchez AJ, Ksiazek TG, Gagro A, Rabatić S, Zgorelec R, Avsić-Zupanc T, Beus I, Dekarić D, 2002. Characteristics of Puumala and Dobrava infections in Croatia. *J Med Virol* 66: 542–551.
 12. Scharninghausen JJ, Pfeffer M, Meyer H, Davis DS, Honeycutt RL, Faule M, 2002. Genetic evidence for Tula virus in *Microtus arvalis* and *Microtus agrestis* populations in Croatia. *Vector Borne Zoonotic Dis* 2: 19–27.
 13. Cvetko L, Markotić A, Plyusnina A, Margaletić J, Miletic-Medved M, Turk N, Milas Z, Avsić-Zupanc T, Plyusnin A, 2005. Puumala virus in Croatia in the 2002 HFRS outbreak. *J Med Virol* 77: 290–294.
 14. Plyusnina A, Krajinović LC, Margaletić J, Niemimaa J, Nemirov K, Lundkvist A, Markotić A, Miletic-Medved M, Avsić-Zupanc T, Henttonen H, Plyusnin A, 2011. Genetic evidence for the presence of two distinct hantaviruses associated with *Apodemus* mice in Croatia and analysis of local strains. *J Med Virol* 83: 108–114.
 15. McFadden G, 2005. Poxvirus tropism. *Nat Rev Microbiol* 3: 201–213.
 16. Tagliapietra V, Rosa R, Hauffe HC, Laakkonen J, Voutilainen L, Vapalahti O, Vaheri A, Henttonen H, Rizzoli A, 2009. Spatial and temporal dynamics of lymphocytic choriomeningitis virus in wild rodents, northern Italy. *Emerg Infect Dis* 15: 1019–1025.
 17. Knap N, Korva M, Dolinsek V, Sekirnik M, Trilar T, Avsić-Zupanc T, 2012. Patterns of tick-borne encephalitis virus infection in rodents in Slovenia. *Vector Borne Zoonotic Dis* 12: 236–242.
 18. Niethamer J, Krapp F, 1982. Handbuch der Säugetiere Europas: Nagetiere 2/1. *Akad Verlag Wiesbaden* 2: 51–491.
 19. Dekonenko A, Yakimenko V, Ivanov A, Morozov V, Nikitin P, Khasanova S, Dzagurova T, Tkachenko E, Schmaljohn C, 2003. Genetic similarity of Puumala viruses found in Finland and western Siberia and of the mitochondrial DNA of their rodent hosts suggests a common evolutionary origin. *Infect Genet Evol* 3: 245–257.
 20. Essbauer S, Schmidt J, Conraths FJ, Friedrich R, Koch J, Hautmann W, Pfeffer M, Wolfel R, Finke J, Dobler G, Ulrich R, 2006. A new Puumala hantavirus subtype in rodents associated with an outbreak of Nephropathia epidemica in south-east Germany in 2004. *Epidemiol Infect* 134: 1333–1344.
 21. Sikes RS, Gannon WL, 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *J Mammal* 92: 235–253.
 22. Tokarz R, Kapoor V, Samuel JE, Bouyer DH, Briese T, Lipkin WI, 2009. Detection of tick-borne pathogens by MassTag polymerase chain reaction. *Vector Borne Zoonotic Dis* 9: 147–152.
 23. Tokarz R, Jain K, Bennett A, Briese T, Lipkin WI, 2010. Assessment of polymicrobial infections in ticks in New York State. *Vector Borne Zoonotic Dis* 10: 217–221.
 24. Bowen MD, Gelbmann W, Ksiazek TG, Nichol ST, Nowotny N, 1997. Puumala virus and two genetic variants of Tula virus are present in Austrian rodents. *J Med Virol* 53: 174–181.
 25. Chu YK, Jennings GB, Schmaljohn CS, 1995. A vaccinia virus-vector Hantaan virus vaccine protects hamsters from challenge with Hantaan and Seoul viruses but not Puumala virus. *J Virol* 69: 6417–6423.
 26. Avsić-Zupanc T, Nemirov K, Petrovec M, Trilar T, Poljak M, Vaheri A, Plyusnin A, 2000. Genetic analysis of wild-type Dobrava hantavirus in Slovenia: co-existence of two distinct genetic lineages within the same natural focus. *J Gen Virol* 81: 1747–1755.
 27. Merien F, Portnoi D, Bourhy P, Charavay F, Berlioz-Arthaud A, Baranton G, 2005. A rapid and quantitative method for the detection of *Leptospira* species in human leptospirosis. *FEMS Microbiol Lett* 249: 139–147.
 28. Victoria B, Ahmed A, Zuerner RL, Ahmed N, Bulach DM, Quinteiro J, Hartskeerl RA, 2008. Conservation of the *SIO-spc-alpha* locus within otherwise highly plastic genomes provides phylogenetic insight into the genus *Leptospira*. *PLoS One* 3: e2752.
 29. Briese T, Palacios G, Kokoris M, Jabado O, Liu Z, Renwick N, Kapoor V, Casas I, Pozo F, Limberger R, Perez-Brena P, Ju J, Lipkin WI, 2005. Diagnostic system for rapid and sensitive differential detection of pathogens. *Emerg Infect Dis* 11: 310–313.
 30. Tamura K, Stecher G, Peterson D, Filipiński A, Kumar S, 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* 30: 2725–2729.
 31. Gravekamp C, Van de Kemp H, Franzen M, Carrington D, Schoone GJ, Van Eys GJ, Everard CO, Hartskeerl RA, Terpstra WJ, 1993. Detection of seven species of pathogenic leptospires by PCR using two sets of primers. *J Gen Microbiol* 139: 1691–1700.
 32. Stritof Majetić Z, Galloway R, Ruzić Sabljic E, Milas Z, Mojčec Perko V, Habus J, Margaletić J, Pernar R, Turk N, 2014. Epizootiological survey of small mammals as *Leptospira* spp. reservoirs in eastern Croatia. *Acta Trop* 131: 111–116.
 33. Turk N, Milas Z, Margaletić J, Staresina V, Slavica A, Riquelme-Sertour N, Bellenger E, Baranton G, Postić D, 2003. Molecular characterization of *Leptospira* spp. strains isolated from small rodents in Croatia. *Epidemiol Infect* 130: 159–166.
 34. Turk N, Milas Z, Margaletić J, Turk R, Barbic L, Konjević D, Perić S, Stritof Z, Staresina V, 2008. The role of fat dromouse (*Glis glis* L.) as reservoir host for spirochete *Borrelia burgdorferi* sensu lato in the region of Gorski Kotar, Croatia. *Eur J Wildl Res* 54: 117–121.
 35. Tea A, Alexiou-Daniel S, Papoutsis A, Papa A, Antoniadis A, 2004. Bartonella species isolated from rodents, Greece. *Emerg Infect Dis* 10: 963–964.
 36. La Scola B, Zeaiter Z, Khamis A, Raoult D, 2003. Gene-sequence-based criteria for species definition in bacteriology: the Bartonella paradigm. *Trends Microbiol* 11: 318–321.
 37. Buffet JP, Kosoy M, Vayssier-Taussat M, 2013. Natural history of Bartonella-infecting rodents in light of new knowledge on genomics, diversity and evolution. *Future Microbiol* 8: 1117–1128.
 38. Welc-Faleciak R, Paziewska A, Bajer A, Behnke JM, Sinski E, 2008. Bartonella spp. infection in rodents from different habitats in the Mazury Lake District, northeast Poland. *Vector Borne Zoonotic Dis* 8: 467–474.
 39. Birtles RJ, Raoult D, 1996. Comparison of partial citrate synthase gene (*gltA*) sequences for phylogenetic analysis of Bartonella species. *Int J Syst Bacteriol* 46: 891–897.
 40. Beck R, Vojta L, Curković S, Mrljak V, Margaletić J, Habrun B, 2011. Molecular survey of Babesia microti in wild rodents in central Croatia. *Vector Borne Zoonotic Dis* 11: 81–83.
 41. Tadin A, Turk N, Korva M, Margaletić J, Beck R, Vucelja M, Habus J, Svoboda P, Zupanc TA, Henttonen H, Markotić A, 2012. Multiple co-infections of rodents with hantaviruses, Leptospira, and Babesia in Croatia. *Vector Borne Zoonotic Dis* 12: 388–392.
 42. Cvetko L, Turk N, Markotić A, Milas Z, Margaletić J, Miletic-Medved M, Plyusnin A, Baranton G, Postić D, Avsić-Zupanc T, 2006. Short report: dual infections with Puumala virus and Leptospira interrogans serovar Lora in a bank vole (Clethrionomys glareolus). *Am J Trop Med Hyg* 74: 612–614.
 43. Sunil-Chandra NP, Clement J, Maes PDE, Silva H, Van Esbroeck M, Van Ranst M, 2015. Concomitant leptospirosis-hantavirus co-infection in acute patients hospitalized in Sri Lanka: implications for a potentially worldwide underestimated problem. *Epidemiol Infect* 143: 2081–2093.
 44. Burri C, Schumann O, Schumann C, Gern L, 2014. Are Apodemus spp. mice and Myodes glareolus reservoirs for Borrelia

- miyamotoi*, *Candidatus* Neorhlichia mikurensis, *Rickettsia helvetica*, *R. monacensis* and *Anaplasma phagocytophilum*? *Ticks Tick Borne Dis* 5: 245–251.
45. Fukunaga M, Takahashi Y, Tsuruta Y, Matsushita O, Ralph D, McClelland M, Nakao M, 1995. Genetic and phenotypic analysis of *Borrelia miyamotoi* sp. nov., isolated from the ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in Japan. *Int J Syst Bacteriol* 45: 804–810.
46. Platonov AE, Karan LS, Kolyasnikova NM, Makhneva NA, Toporkova MG, Maleev VV, Fish D, Krause PJ, 2011. Humans infected with relapsing fever spirochete *Borrelia miyamotoi*, Russia. *Emerg Infect Dis* 17: 1816–1823.
47. Hovius JW, de Wever B, Sohne M, Brouwer MC, Coumou J, Wagemakers A, Oei A, Knol H, Narasimhan S, Hodiamont CJ, Jahfari S, Pals ST, Horlings HM, Fikrig E, Sprong H, van Oers MH, 2013. A case of meningoencephalitis by the relapsing fever spirochaete *Borrelia miyamotoi* in Europe. *Lancet* 382: 658.
48. Gugliotta JL, Goethert HK, Berardi VP, Telford SR 3rd, 2013. Meningoencephalitis from *Borrelia miyamotoi* in an immunocompromised patient. *N Engl J Med* 368: 240–245.
49. Bown KJ, Begon M, Bennett M, Woldehiwet Z, Ogden NH, 2003. Seasonal dynamics of *Anaplasma phagocytophila* in a rodent-tick (*Ixodes trianguliceps*) system, United Kingdom. *Emerg Infect Dis* 9: 63–70.